



# 129

## Origin

Developed by Dunn at Columbia University in 1928 from a cross of coat color stocks and a chinchilla (*Tyr<sup>c-ct</sup>*) stock from Castle. Strain 129 has a common origin with strain 101. In 1945, this strain was passed to Russell at the Jackson Laboratory, Bar Harbor. This stock was maintained segregating at the *Tyr* locus, such that the coat color of the mice were either albino or light chinchilla. In 1947, to Hunt at Michigan State and in 1948 again introduced at the Jackson Laboratory (129/ReJ) after a forest fire.

## 129P2/OlaHsd

In 1978, to GD Searle, High Wycomb. From GD Searle to Olac, which became Harlan, in 1979. Harlan became Envigo in 2015.

## 129S2/SvHsd

In 1953, to LC Stevens at the Jackson Laboratory. Stevens initially studied the genetic basis of testicular teratomas in mice of the 129 parental strain. To determine the cellular origin of teratomas, he performed outcrosses to introduce onto 129 background mutations that affect gonads or germ cells. This led to the "Steel" and "Ter" substrains. The Steel substrains originated from an outcross to C3H-*Mgf<sup>st-J</sup>*, followed by 12-14 generations of backcrossing to 129/Sv. The resulting mice are pigmented white-bellied agouti. The steel-J mutation increased the teratoma incidence to 10%. From Glaxo-Wellcome to Harlan Laboratories. Harlan became Envigo in 2015

## Nomenclature

There is substantial genetic variation among substrains of strain 129 (Simpson *et al*, 1997; Threadgill *et al*, 1997). Therefore, the Committee on Standardized Genetic Nomenclature for Mice has approved a new nomenclature for strain 129 (Festing *et al*, 1999). The new nomenclature is based on the substrains identified and defined in terms of microsatellite markers by Simpson *et al* (1997). A

letter and a number have been introduced in front of the slash that will identify each of the substrains. The letter is either P, S, T or X indicating whether it is a 'Parental', 'Steel', 'Ter' (susceptible to teratomas) or genetically-contaminated "X" substrain, respectively.

The major parental lineages include:

- 129P - derived from the original parent strain
- 129S - derived from a congenic strain made by outcrossing the *steel* mutation
- 129T - derived from the 129 congenic that originally carries the teratoma mutation
- 129X - derived from a genetically-contaminated "X" substrain

Therefore the 129/OlaHsd strain is renamed into 129P2/OlaHsd and 129/SvHsd is renamed into 129S2/SvHsd.

The choice of a particular substrain may be of critical importance for a particular project.

## Research applications

Transgenesis, testicular teratomas.

## Characteristics

Strain 129 is useful for ovary transplant and ova transfer studies. It is best known for the high incidence of testicular teratomas, but more recently it has been the most widely-used strain in the production of targeted mutations due to the availability of several lines of embryonic stem cells.

## Anatomy

Large brain/body weight ratio, small spinal cord (Roderick *et al*, 1973). Small forebrain volume and neocortex (Wimer *et al*, 1969). A large proportion of 129P2/OlaHsd mice has major shunts between the hepatic portal system and the vena cava, allowing the passage of microspheres up to 50  $\mu$ m in diameter.

These shunts are associated with resistance to *Shistosoma japonicum* cercariae (Coulson and Wilson, 1989). High retinal ganglion cell number in 129P3/J substrain (Williams *et al*, 1996). Absence of corpus callosum in about 70% of mice of the 129P3/J substrain. This may be related to retarded formation of the hippocampal commissure in this strain and in BALB/c mice (Livy and Wahlsten, 1997). High bone density of femur in 129P3/J substrain (Beamer *et al*, 1996).

### Behavior

Low avoidance conditionability (Royce, 1972). Low shock-avoidance learning (Royce *et al*, 1971). Low preference for sweet tasting substances (saccharin, sucrose, dulcin and acesulfame, averaged 129S substrain) (Lush 1988). Prefers moderate concentrations of saline (contrast C57BL/6) (Beauchamp and Fisher, 1993; Gannon and Contreras, 1995). Normal sensitivity to provoked ataxia by diazepam (Crabbe *et al*, 1998).

### Drugs

Resistant to skin ulceration by DMBA (Thomas *et al*, 1973.). Resistant to induction of subcutaneous tumors by 3-methylcholanthrene (Kouri *et al*, 1973.). Resistant to X-irradiation (Roderick, 1963; Storer, 1966). Females have long sleeping time under hexobarbital anesthetic (Lovell, 1976). Resistant to toxic effects of isoniazid (Taylor, 1976). Insensitive (eosinophil response) to cortisone acetate (Wragg and Speirs, 1952). Sensitive uterine response to estrogens (Chai and Dickie, 1966; Drasher, 1965). Susceptible to ozone-induced decreases of tracheal potential (Takahashi *et al*, 1995).

### Genetics

#### 129P2/OlaHsd:

Coat color genes -  $A^w$ ,  $B$ ,  $c^{ch}$ ,  $D$ ,  $p$ :  
pink-eyed chinchilla.

Histocompatibility -  $H-1^b$ ,  $H-2^b$ ,  $H-3^b$ .

#### 129S2/SvHsd:

Coat color genes -  $A^w$ ,  $C$ ,  $P$ :  
white-bellied agouti.

Substrain 129P2/OlaHsd differs from 129S2/SvHsd at many loci, particularly those on chromosome seven, which has the albino and pink-eyed dilute loci.

Most substrains carry the white-bellied agouti gene  $A^w$  though only a subset have the agouti pattern as many carry albino or chinchilla and/or the pink-eyed dilution gene,  $p$ , which is derived from Asian mice of the *Mus musculus* type (Brilliant *et al*, 1994).

This strain carries the *Mus musculus musculus* Y-chromosome, while others have the *M. m. domesticus* type (Nishioka, 1987).

### Immunology

High lymphocyte phytohemagglutinin response (Heiniger *et al*, 1975). Responder to synthetic polypeptide (Glu<sup>57</sup>, Lys<sup>38</sup>, Ala<sup>5</sup>) (Pinchuck and Maurer, 1965). Erythrocytes have a high agglutinin ability (Rubinstein *et al*, 1974). High responder to Dextran (Blomberg *et al*, 1972). Experimental systemic lupus

erythematosus including severe ocular changes and blepharitis can be induced by injection of human monoclonal anti-DNA antibodies (Chan *et al*, 1995).

### Infection

Carries no detectable endogenous ecotropic MuLV DNA sequences (Jenkins *et al*, 1982). Intermediate sensitive to *Yersinia enterocolitica*, while C57BL/6 is resistant and BALB/c is sensitive (Bohn *et al*, 1998).

### Life-span and spontaneous disease

Low overall tumor incidence (7% in males, 21% in females), including lymphoma 2% in males and 7% in females, soft tissue sarcomas 2% in males and 1% in females and benign tumors 2% in males and 3% in females (Smith *et al*, 1973). Lung tumors 4-46% (Festing and Blackmore, 1971). Testicular teratomas about 1% in most substrains, but 30% in the terSv substrain (Stevens, 1973). Incidence of teratomas increased in p53-deficient mice (Harvey *et al*, 1993). The Ter gene has been mapped to chromosome 18 (Asada *et al*, 1994). High incidence of urinary calculi (Russell and Meier, 1966). A review of the life span of aging mice has been described by Myers, (1978). Median life-span 22.9 months in 129P3/J males and 21.6 months in 129P3/J females (Storer, 1966). Median life-span 27.3 months in 129P3/J males and 24.3 months in 129P3/J females (Smith *et al*, 1973). Median life-span 23.3 months in 129/RrJLac males and 22.3 months in 129/RrJLac females (Festing and Blackmore, 1971).

### Normal physiology and biochemistry

High plasma cholesterol at 12 and 24 weeks (Weibust, 1973). Low plasma triglyceride levels (Jiao *et al*, 1990). High Na/K ratio in erythrocytes and plasma (Waymouth, 1973). High serum ceruloplasmin levels in males but low levels in females (Meier and MacPike, 1968). High plasma cholinesterase activity (Angel *et al*, 1967). Low mean heart rate but high mean heart adaptation rate (Blizard and Welty, 1971). High cell turnover in 129P3/J substrain as estimated by rapid clearance of DNA-bound radioactivity (Heiniger *et al*, 1972). Venous blood has a high pH (Bernstein, 1966). High hepatic microsomal coumarin hydroxylase activity in females (Van Iersel *et al*, 1994). High levels of apoA-IV messenger RNA in liver compared with C57BL/6 (Reue *et al*, 1993). Has defective secretory group II phospholipase A2 gene (confer strains C57BL/6 and B10.RIII) (Kennedy *et al*, 1995).

### Miscellaneous

Recommended host for transplantable tumor hemangi endothelioma BW6473 (Kaliss, 1972). The E14 clone of embryonic stem cells was derived from mice of 129P2 substrain. (Hooper *et al*, 1987; Papaioanou and Johnson, 1993). Substrain 129S2 is widely used in gene targeting experiments because most of the available embryonic stem cell lines have been developed from strain 129S2. The D3 clone of embryonic stem cells was derived from mice of this substrain (Doetschman *et al*, 1985; Papaioanou and Johnson, 1993). Characteristics of the 129 strain have been described by Festing (1997) and Lyon (1996).

### Production

Poor breeding performance, colony output 0.8 young/female/wk, litter size at weaning 4.5 (Festing, 1976)

## References

- Angel CR, Mahin DT, Farris RD, Woodward KT (1967) Heritability of plasma cholinesterase activity in inbred mouse strains. *Science* 156, 529-530.
- Asada Y, Varnum DS, Frankel WN, Nadeau JH (1994) A mutation in the Ter gene causing increased susceptibility to testicular teratomas maps to mouse chromosome 18. *Nature Genet.* 6, 363-368.
- Beamer WG, Donahue LR, Rosen CJ, Baylink DJ (1996) Genetic-variability in adult bone-density among inbred strains of mice. *Bone* 18, 397-403.
- Beauchamp GK, Fisher AS (1993) Strain differences in consumption of saline solutions by mice. *Physiol. Behav.* 54, 179-184.
- Bernstein SE (1966) Physiological characteristics. In: *Biology of the Laboratory Mouse*. 2nd ed. (Green EL, ed). New York: McGraw-Hill, pp. 337-350.
- Blizard DA, Welty R (1971) Cardiac activity in the mouse: strain differences. *J. Comp. Physiol. Psychol.* 77, 337-344.
- Blomberg B, Geckeler WR, Weigert M (1972) Genetics of the antibody response to Dextran in mice. *Science* 177, 178-180.
- Bohn E, Schmitt E, Bielefeldt C, Noll A, Schulte R, Autenrieth IB (1998) Ambiguous role of interleukin-12 in *Yersinia enterocolitica* infection in susceptible and resistant mouse strains. *Infect. Immun.* 66, 2213-2220.
- Brilliant MH, Ching A, Nakatsu Y, Eicher EM (1994) The original pink-eyed dilution mutation (p) arose in asiatic mice: Implications for the H4 minor histocompatibility antigen, *Myod1* regulation and the origin of inbred strains. *Genetics* 138, 203-211.
- Chai CK, Dickie MM (1966) Endocrine variations. In *Biology of the laboratory mouse*. 2nd ed. (Green EL, ed), pp. 387-403. McGraw-Hill, New York.
- Chan CC, Gery I, Kohn LD, Nussenblatt RB, Mozes E, Singer DS (1995) Periocular inflammation in mice with experimental systemic lupus erythematosus: A new experimental blepharitis and its modulation. *J. Immunol.* 154, 4830-4835.
- Coulson PS, Wilson RA (1989) Portal shunting and resistance to *Shistosoma mansoni* in 129 strain mice. *Parasitology* 99, 383-389.
- Crabbe JC et al (1998) Genetic determinants of sensitivity to diazepam in inbred mice. *Behav. Neurosci.* 112, 668-677.
- Doetschman TC, Eistetter H, Katz M, Schmidt W, Kemler M (1985) The *in vitro* development of blastocyst-derived embryonic stem cell lines: formation of visceral yolk sac, blood island and myocardium. *J. Embryol. Exp. Morph.* 87, 27-45.
- Drasher ML (1965) Strain differences in the response of the mouse uterus to estrogens. *J. Hered.* 46, 190-192.
- Festing MFW (1976) Effects of marginal malnutrition on the breeding of inbred and F1 hybrid mice – a diallel study. In: *The Laboratory Animal in the Study of Reproduction*. (Antikatzides T, Ericksen S, Spiegel A, eds). Stuttgart: Gustav Fischer, pp 99-114.
- Festing MFW (1997) Inbred Strains of mice. *Mouse genome* 95, 519-686.
- Festing MFW, Blackmore DK (1971) Life span of specified-pathogen-free (MRC category 4) mice and rats. *Lab. Anim.* 5, 179-192.
- Festing MFW, Simpson EM, Davisson MT, Mobraaten LE (1999) Revised nomenclature for strain 129 mice. *Mammalian Genome* 10, 836.
- Gannon KS, Contreras RJ (1995) Sodium intake linked to amiloride-sensitive gustatory transduction in C57BL/6J and 129/J mice. *Physiol. Behav.* 57, 231-239.
- Harvey M, McArthur MJ, Montgomery CA, Bradley A, Donehower LA (1993) Genetic background alters the spectrum of tumors that develop in p53-deficient mice. *FASEB* 7, 938-943.
- Heiniger HJ, Chen HW, Meier H, Taylor BA, Commerford LS (1972) Studies on the genetic control of cell proliferation. 1. Clearance of DNA-bound radioactivity in 19 inbred strains and hybrid mice. *Life Sci.* 11, 87-98.
- Heiniger HJ, Taylor BA, Hards EJ, Meier H (1975) Heritability of the phytohemagglutinin responsiveness of lymphocytes and its relationship to leukemogenesis. *Cancer Res.* 35, 825-831.
- Hooper M, Hardy K, Handyside A, Hunter S, Monk M (1987) HPRT-deficient (Lesch-Nyhan) mouse embryos derived from germline colonization by cultured cell. *Nature* 326, 292-295.
- Jenkins NA, Copeland NG, Taylor BA, Lee BK (1982) Organization, distribution and stability of endogenous ecotropic murine leukemia virus DNA sequences in chromosomes of *Mus musculus*. *J. Virol.* 43, 26-36.
- Jiao S, Cole TG, Kitchens R, Pfleger B, Schonfeld G (1990) Genetic heterogeneity of lipoproteins in inbred strains of mice: analysis by gel-permeation chromatography. *Metabolism* 39, 155-160.
- Kaliss N (1972) Transplanted tumors. *Jax Notes* 410. Jackson Laboratory, Bar Harbor, Maine.
- Kalter H (1968) Sporadic congenital malformations of newborn inbred mice. *Teratology* 1, 193-200.
- Kennedy BP, Payette P, Mudgett J, Vadas P, Pruzanski W, Kwan M, Tang C, Rancourt DE, Cromlish WA (1995) A natural disruption of the secretory group II phospholipase A2 gene in inbred mouse strains. *J. Biol. Chem.* 270, 22378-22385.
- Kouri RE, Salerno RA, Whitmire CE (1973) Relationships between arylhydrocarbon hydroxylase inducibility and sensitivity to chemically induced subcutaneous sarcomas in various strains of mice. *J. Natl. Cancer Inst.* 50, 363-368.
- Livy DJ, Wahlsten D (1997) Retarded formation of the hippocampal commissure in embryos from mouse strains lacking a corpus callosum. *Hippocampus* 7, 2-14.
- Lovell DP (1976) Hexobarbital sleeping time in mice. Unpublished data.
- Lush IM (1988) The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet. Res.* 53, 95-99.
- Lyon MF, Rastan S, Brown SDM (1996) Genetic variants and strains of the laboratory mouse. 2 Volumes. Oxford, New York, Tokyo: Oxford University Press.
- Meier H, MacPike AD (1968) Levels and heritability of serum ceruloplasmin activity in inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 128, 1185-1190.
- Myers DD (1978) Review of disease patterns and life span in aging mice: Genetic and environmental interactions. *Birth defects: Original article series* 14, 43-51.
- Nishioka Y. (1987) Y-chromosomal DNA polymorphism in mouse inbred strains. *Genet. Res.* 50, 69-72.
- Papaioanou V, Johnson R (1993) Production of chimeras and genetically defined offspring from targeted ES cells. In: *Gene targeting*. (Joyner AL, ed). Oxford, New York: Oxford University Press, pp 107-146.
- Pinchuck P, Maurer PH (1965) Antigenicity of polypeptides (poly alpha amino acids). XVI. Genetic control of immunogenicity of synthetic polypeptides in mice. *J. Exp. Med.* 122, 673-679.
- Reue K, Purcell-Huynh DA, Leete TH, Doolittle MH, Durstenfeld A, Lusa AJ (1993) Genetic variation in mouse apolipoprotein A-IV expression is determined pre- and post-transcriptionally. *J. Lipid Res.* 34, 893-903.
- Roderick TH (1963) The response of twenty-seven inbred strains of mice to daily doses of whole-body X-irradiation. *Radiation Res.* 20, 631-639.
- Roderick TH, Wimer RE, Wimer CC, Schwartzkroin PA (1973) Genetic and phenotypic variation in weight of brain and spinal cord between inbred strains of mice. *Brain Res.* 64, 345-353.
- Royce JR (1972) Avoidance conditioning in nine strains of inbred mice using optimal stimulus parameters. *Behav. Genet.* 2, 107-110.
- Royce JR, Yeudall LT, Poley W (1971) Diallel analysis of avoidance conditioning in inbred strains of mice. *J. Comp. Physiol. Psychol.* 76, 353-358.
- Rubinstein P, Liu N, Strenn EW, Decary F (1974) Electrophoretic mobility and agglutinability of red blood cells: a 'new' polymorphism in mice. *J. Exp. Med.* 139, 313-322.
- Russell ES, Meier H (1966) Constitutional diseases. In: *Biology of the Laboratory Mouse*. 2nd ed. (Green EL, ed). pp. 571-587. McGraw-Hill, New York.
- Simpson EM, Linder CC, Sargent EE, Davisson MT, Mobraaten LE and Sharp JJ (1997) Genetic variation among substrains and its importance for targeted mutagenesis in mice. *Nature Genet.* 16, 19-27.
- Smith GS, Walford RL, Mickey RM (1973) Lifespan and incidence of cancer and other diseases in selected long-lived inbred mice and their F1 hybrids. *J. Natl. Cancer Inst.* 50, 1195-1213.
- Stevens LC (1973) A new inbred subline of mice (129/ter, Sv) with a high incidence of spontaneous congenital testicular teratomas. *J. Natl. Cancer Inst.* 50, 235-242.
- Storer JB (1966) Longevity and gross pathology at death in 22 inbred strains of mice. *J. Gerontol.* 21, 404-409.
- Takahashi M, Kleiberger SR, Croxton TL (1995) Genetic control of susceptibility to ozone-induced changes in mouse tracheal electrophysiology. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 269, L6-L10.
- Taylor BA (1976) Genetic analysis of susceptibility to isoniazid-induced seizures in mice. *Genetics* 83, 373-377.
- Thomas PE, Hutton JJ, Taylor BA (1973) Genetic relationship between aryl hydrocarbon hydroxylase inducibility and chemical carcinogen induced skin ulceration in mice. *Genetics* 74, 655-659.
- Threadgill DW, Yee D, Matin A, Nadeau JH, Magnuson T (1997) Genealogy of the 129 inbred strains: 129/SvJ is a contaminated inbred strain. *Mamm. Genome* 8, 390-393.
- Van Iersel M, Walters DG, Price RJ, Lovell DP and Lake BG (1994) Sex and strain differences in mouse hepatic microsomal coumarin 7-hydroxylase activity. *Food and Chemical Toxicology* 32, 387-390.
- Waymouth C (1973) Erythrocyte sodium and potassium levels in normal and anemia mice. *Comp. Biochem. Physiol.* 44A, 751-766.
- Weibust RS (1973) Inheritance of plasma cholesterol levels in mice. *Genetics* 73, 303-312.
- Williams RW, Strom RC, Rice DS, Goldowitz D (1996) Genetic and environmental-control of variation in retinal ganglion-cell number in mice. *Journal of Neuroscience* 16, 7193-7205.
- Wimer RE, Wimer CC, Roderick TH (1969) Genetic variability in forebrain structures between inbred strains of mice. *Brain Res.* 16, 257-264.
- Wragg LE, Speirs RS (1952) Strain and sex differences in response of inbred mice to adrenal cortical hormones. *Proc. Soc. Exp. Biol. Med.* 80, 680-684.

## Contact us

North America 800.793.7287 EU and Asia [envigo.com/contactus](http://envigo.com/contactus) [info@envigo.com](mailto:info@envigo.com)



Envigo RMS Division, 8520 Allison Pointe Blvd., Suite 400, Indianapolis, IN 46250, United States

© 2016 Envigo.

RMS-0216-EU-01-PS-185