



+++
ENVIGO

Research Models
and Services

Inbred Mice

C57BL/6

Origin

Developed in 1921 by Little from brother - sister pair (female 57 x male 52) of Miss Abby Lathrop's stock. The same cross gave rise to strains C57L and C57BR. Female 58 mated with the same male gave rise to strain C58. Strains 6 and 10 separated prior to 1937. In 1946, to the Jackson Laboratory, Bar Harbor.

C57BL/6JOlaHsd

In 1974, from the Jackson Laboratory to Laboratory Animals Centre, Carshalton. To OLAC (now Envigo) in 1983. In 1997 to Harlan Nederland (now Envigo).

C57BL/6NHsd

In 1974, from the Jackson Laboratory to the National Institutes of Health, Bethesda, Maryland. Harlan Sprague Dawley, Inc., derived the strain from this breeding nucleus. Harlan became Envigo in 2015.

C57BL/6JRccHsd

In 1973, from the Jackson Laboratory to the Biological Research Laboratories-RCC Ltd. Füllinsdorf, Switzerland. In 2005, Harlan obtained a breeding nucleus after acquisition of RCC Ltd. Harlan became Envigo in 2015.

Research applications

Behavior, learning, atherosclerosis, metabolism, alcohol preference anatomy, irradiation, carcinogenesis, immunology, infections.

Characteristics

The C57BL is easily the most widely used of all inbred strain. Used as a genetic background for many mutants e.g. obese, diabetes and beige. This is a long-lived strain with few tumors, some spontaneous congenital abnormalities.

Anatomy

Small kidney/body weight ratio (Schlager, 1968). Large thyroid (Mendoza *et al*, 1967). High total leukocyte count, low erythrocyte count (Russell *et al*, 1951). Small hippocampus (Wimer *et al*, 1969). Accessory spleens in about 32% of mice and low number of Peyer's patches (Hummel *et al*, 1966). Higher bone mass than A/J (Kaye and Kusy, 1995). Hematopoietic stem cell pool 11-fold lower than in DBA/2. This is largely due to loci on chromosome 1 (Muller-Sieburg and Riblet, 1996). Less susceptible to the development of micronuclei than BALB/c following treatment with clastogenic base analogues and nucleosides (Sato *et al*, 1993). High level of spontaneous sister chromatid exchange (Nishi *et al*, 1993). A detailed staging of these mice between gestation days 11 and 13 (Theiler's stages 18 and 21) has been published by Miyake *et al*, (1996a). Low bone density of femur (Beamer *et al*, 1996; 2001). The timing of onset and duration of condensation and onset of matrix formation of first arch cartilages has been described by Miyake *et al* (1996a). A detailed staging table to facilitate study of cranial skeletal development every two hrs between days 11 and 13 of gestation has also been described (Miyake *et al*, 1996b).

Behavior

High alcohol (ethanol) preference (Fuller, 1964; Rodgers, 1966). The mean maximally preferred concentrations of ethanol were 17,9% for C57BL/6 and 6,8% for ICR mice. The consumption of ethanol represents a preferred source of calories for the C57BL/6 mouse (McMillen *et al*, 1998).

Achieve blood alcohol levels of 60 mg% when access to alcohol is restricted to 60 min per day (Le *et al*, 1994). Alcohol preference may be associated with strain differences in mesolimbic enkephalin gene expression (Ng *et al*, 1996). A quasi-congenic QTL introgression strain carrying a low alcohol consumption gene from BALB/c has lower voluntary alcohol consumption than C57BL/6, with 96% of loci in common (Vadasz *et al*, 1996).

Low severity of ethanol withdrawal symptoms compared with DBA/2, possibly associated with differences in neuroactive steroid sensitivity (Finn *et al*, 1997). Alcohol preference is due to at least two recessive quantitative trait loci that are sex-restricted in expression (Melo *et al*, 1996). Low 'emotionality', high open-field exploration (Thompson, 1953). High spontaneous locomotor activity (Nikulina *et al*, 1991). Short time of immobility in a forced swimming test (Nikulina *et al*, 1991). Low shock-avoidance learning (Bovet *et al*, 1966). Low shuttle-box avoidance, high wheel activity (Messeri *et al*, 1972). Rapid shock-avoidance learning and slow extinction (Schlesinger and Wimer, 1967). High shock-avoidance learning (Wahlsten, 1973). High radial-arm maze learning (Ammassari-Teule *et al*, 1993). High locomotor activity (Davis and King, 1967). High locomotor activity when grouped and single (Davis *et al*, 1967). Resistant to audiogenic seizures (Fuller and Sjursen, 1967). Relatively insensitive to the primary odorant isovaleric acid (contrast seven other strains) and may provide an animal model of specific anosmia (Wysocki *et al*, 1977). Low balsa-wood gnawing activity (Fawcington and Festing, 1980). High preference for sweet tasting substances (saccharin, sucrose, dulcin and acesulfame, averaged) (Lush 1988). Rejects saline at moderate concentrations (contrast 129) (Beauchamp and Fisher, 1993, Gannon and Contreras, 1995). Feed restriction for nine days failed to cause stereotypic cage cover climbing (contrast DBA/2) (Cabib and Bonaventura, 1997). In C57BL/6 mice self-grooming and allo-grooming is observed (Miltzer and Wecker, 1986)

Drugs

Susceptible to skin ulceration by DMBA (Thomas *et al*, 1973). Susceptible to induction of subcutaneous tumors by 3-methylcholanthrene (Kouri *et al*, 1973; Whitmire *et al*, 1971). High incidence of lymphomas after methylcholanthrene administration by gavage (Akamatsu and Barton, 1974). Susceptible to toxic effects of DMBA (Schmid *et al*, 1966). Pre-treatment with beta-naphthoflavone 48 hr. before administration of N-nitrosoethylurea (ENU), once weekly for four weeks caused a significant doubling in the number of lung tumor bearers (contrast 4 strains) (Anderson *et al*, 1990). Phenobarbitone in the diet to give an intake of 85 mg/kg per day resulted in 4% of animals developing basophilic nodules by 91 weeks of age (contrast 70% in C3H/He), but no increase in liver carcinomas (Evans *et al*, 1992). However, there was a two-fold lower level of DNA synthesis in C57BL/6 mice relative to C3H mice after partial hepatectomy, though partial hepatectomy is a tumor promoter in C57BL/6 but not in C3H mice (Bennett *et al*, 1995). Sensitive to teratogenic effects of acetazolamide (Green *et al*, 1973). Resistant to teratogenic effect (cleft palate) by cortisone acetate (Kalter 1981). Hepatic epoxide hydrase activity induced by pentobarbital i.p. (Oesch *et al*, 1973).

Resistant to teratogenic effects of cortisone acetate (Dostal and Jelinek, 1973). Resistant to lethal effects of ozone (Goldstein *et al*, 1973), but susceptible to ozone-induced decreases of tracheal potential (Takahashi *et al*, 1995) and to airway inflammation (contrast C3H/He) (Kleeberger *et al*, 1993). Susceptible to ozone-induced lung

inflammation, which is exacerbated by vitamin A deficiency (Paquette *et al*, 1996). High incidence of convulsions induced by flurothyl (Davis and King, 1967). Susceptible to hyperbaric oxygen (Hill *et al*, 1968). Resistant to chloroform toxicity (Hill *et al*, 1975; Deringer *et al*, 1953). Resistant to toxic effects of isoniazid (Taylor 1976). Sensitive, as judged by eosinophil response, to cortisone acetate (Wragg and Speirs, 1952). High (89%) ovulatory response to three I.U. of PMS in immature mice, but only a 56% response to 7 I.U. No facilitation by exposure to males at these doses (Zarrow *et al*, 1971). High locomotor activity after treatment with σ -amphetamine (Babbini *et al*, 1974). Nicotine increases learning ability (Bovet *et al*, 1966). Resistant to colon carcinogenesis by 1,2-dimethylhydrazine (Evans *et al*, 1977). Low ED50 to behavioral effects of nicotine (Marks *et al*, 1989). High self-selection of nicotine which is inversely correlated with sensitivity to nicotine-induced seizures (Robinson *et al*, 1996). Low bronchial reactivity to methacholine and serotonin (Konno *et al*, 1993). Resistant to daunomycin-induced nephrosis (Kimura *et al*, 1993).

Low neural sensitivity to pentylenetetrazol convulsions (Kosobud *et al*, 1992). Susceptible to biliary tract injury following oral dosing with 500 micrograms of the fungal toxin sporidesmin (Bhathal *et al*, 1990). Low histamine release from peritoneal mast cells induced by compound 48/80, a calcium dependent histamine releaser (Toda *et al*, 1989). Low histamine release from peritoneal mast cells induced by Ca²⁺ ionophore A23187, (contrast BALB/c, C3H/He, DBA/2 etc.) (Toda *et al*, 1989). Carries gene (Tpmt) for low levels of thiopurine methyltransferase activity, catalyzing the S-methylation of 6-mercaptopurine and other heterocyclic and aromaticthiol compounds (like AKR, unlike DBA/2) (Otterness and Weinsilbourn 1987a;b). More sensitive to acute toxic effects of aflatoxin B-1 than strains CBA/J or BALB/c (Almeida *et al*, 1996). Airways hyporeactive to acetylcholine (Zhang *et al*, 1995). High voluntary consumption of morphine in two-bottle choice situation (Belknap *et al*, 1993). Estrogen induces an increase in VLDL and LDL-cholesterol (like C57L, contrast BALB/c and C3H) (Srivastava, 1995). Nine-fold higher ED50 for haloperidol-induced catalepsy than DBA/2, but this is not associated with numbers of cholinergic neurons (Dains *et al*, 1996). Accumulates three to five-fold lower levels of mercury in liver and blood than DBA/2 or A.SW after four weeks exposure to mercuric chloride, but higher levels in spleen following 8-12 weeks of exposure (Griem *et al*, 1997).

Genetics

Coat color genes	- a, B, C, D : black.
Histocompatibility	- H-1 ^c , H-2 ^b , H-3 ^a .
Biochemical markers	- Apoa-1 ^a , Car-2 ^a , Es-1 ^a , Es-2 ^b , Es-3 ^a , Gpd-1 ^a , Gpi-1 ^b , Hba ^a , Hbb ^s , Idh-1 ^a , Ldr-1 ^a , Mod-1 ^b , Mup-1 ^b , Pep-3 ^a , Pgm-1 ^a , Pgm-2 ^a , Trf ^b .

Four major substrains A, GrFa, 6 and 10 appear to be quite similar, and any differences are consistent with what might be expected from the accumulation of new mutations and a small amount of residual heterozygosity, though McClive *et al* (1994) have found that B6 and B10 differ at multiple loci on chromosome 4 including the microsatellite markers D4Mit69, D4Mit71 and D4Mit72. Additional microsatellites, which distinguish between B6 and B10 are given by Slingsby *et al* (1996). Substrains 6 and 10 differ at the *H-9*, *Igh-2* and *Lv* loci. All Envigo C57BL/6 sublines still carry the Nnt (nicotinamide nucleotide transhydrogenase) gene, which is missing in the original C57BL/6J from Jackson Laboratories.

C57BL/6J α Hsd mice lack α -synuclein due to a small deletion of the locus (Specht and Schoepfer, 2001). α -Synuclein belongs to a family of structurally related proteins expressed highly in the brain. However, α -synuclein is not essential for spatial learning tasks (Chen *et al*, 2002). This deletion isn't present in the C57BL/6JRccHsd subline!

Description of the difference between FVB/N and C57BL/6J for 272 microsatellites (Neuhaus *et al*, 1997). A probe designated B6-38 to the pseudoautosomal region of the X and Y chromosome has a characteristic Pst I pattern of fragment sizes which is present only in the C57BL family of strains (Kalcheva *et al*, 1995). C57BL/6 mice carry the *Mus musculus* Y-chromosome, while others have the *M. m. domesticus* type (Nishioka, 1987).

The C57BL/6NHsd subline carries a retinal degeneration 8 mutation - *rd8* (Caspi *et al*, 2012)

Immunology

High susceptibility to induction of amyloid by casein (Willerson *et al*, 1969). Poor immune response to type III pneumococcal polysaccharide (Braley and Freeman, 1971). Poor immune response to synthetic double-stranded RNA (Steinberg *et al*, 1971). Good immune response to cholera A and B antigens (Cerny *et al*, 1971). Resistant to induction of anaphylactic shock by ovalbumin (Tanioka and Esaki, 1971). Rapid rejection of about 76% of male skin isografts by females by 25 days (Gasser and Silvers, 1971). Poor immune response to GAT (random terpolymer of Glu⁶⁰, Ala³⁰, Tyr¹⁰) (9/10) (Dorf *et al*, 1974). Good immune response to *Salmonella senftenberg* and *S. anatum* lipopolysaccharide (Di Pauli, 1972). Non-responder to synthetic polypeptide Glu⁵⁷, Lys³⁸, Ala⁵ (Pinchuck and Maurer, 1965). High sporadic occurrence of natural hemagglutinins to sheep red blood cells (Brooke, 1965). Discriminator between 'H' and 'L' sheep erythrocytes (McCarthy and Dutton, 1975).

Poor immune response to Pro-Gly-Pro-ovalbumin and (Pro⁶⁶, Gly³⁴)_n but good immune response to (Pro-Gly-Pro)_n (Fuchs *et al*, 1974). High PHA-stimulated lymphocyte blastogenic response (Hellman and Fowler, 1972). Erythrocytes have low agglutinability (Rubinstein *et al*, 1974). High immune response to ferritin in B6-*T1a* (Young *et al*, 1976). Low responder to dextran (Blomberg *et al*, 1972). Low responder to *E. coli* β -D-galactosidase, with "memory" developing in absence of antibody formation (De Macario and

Macario 1980). Precipitating and skin sensitising antibodies have slow electrophoretic mobility (Fahey, 1965). Resistant to anaphylactic shock (Treadwell, 1969). Susceptible to induction of autoimmune prostatitis (contrast BALB/c) (Keetch *et al*, 1994). High expression of neutral glycosphingolipid GgOse(4) Cer in concanavalin A stimulated T lymphoblasts (Muthing, 1997). Anti-BPO IgE monoclonal antibody produced potent systemic sensitization sufficient for provocation of lethal shock in most aged (6 to 10 months) mice (Harada *et al*, 1991). Susceptible to immunosuppression of contact hypersensitivity by ultraviolet B light (Noonan and Hoffman, 1994). The potential influence of circadian changes and laboratory routine on some immune parameters has been described by Kolaczowska *et al* (2000).

Infection

Develops a slowly progressing parasitosis ("low responder") after infection with the Cornell strain of *Toxoplasma gondii* (Macario *et al*, 1980). Did not support sustained growth of six strains of *Leishmania mexicana mexicana* (contrast BALB/c) (Monroy-Ostria *et al*, 1994). Resistant to *Leishmania major* (contrast BALB/c) (Laskay *et al*, 1995; Scott *et al*, 1996). Susceptible to *L. major mexicana*, and vaccination against the parasite using liposomes with parasite membrane antigens was effective (cf CBA/Ca but contrast C57BL/10) (Lezama-Dávila, 1997). Susceptible to *Salmonella typhimurium* strain C5 (Robson and Vas, 1972). 100-fold more resistant to *Listeria monocytogenes* than A/J when measured by median lethal dose (Sadarangani *et al*, 1980). This seems to be associated with increased levels of gamma interferon and granulocyte-macrophage colony stimulating factor compared with susceptible A/J mice (Iizawa *et al*, 1993). Resistant to *Mycoplasma fermentans* (Gabridge *et al*, 1972). Resistant to *Mycoplasma pulmonis* infection (Cartner *et al*, 1996). Resistant to infection by *Mycobacterium marinum* (Yamamoto *et al*, 1991). Resistant to infection by liver fluke *Opisthorchis felineus* (Zelentsov, 1974). Resistant to infection with the helminth worm *Angiostrongylus costaricensis* (Ishii and Sano 1989). Relatively susceptible to infection with *Helicobacter felis* (contrast C57BL/6) (Mohammadi *et al*, 1996). Susceptible to infection by *Helicobacter felis* with moderate to severe chronic active gastritis in the body of the stomach, which increased over time (Sakagami *et al*, 1996). *H. felis* induces hypertrophic gastropathy (Fox *et al*, 1996). Highly resistant to the mammary tumor virus which is thought not to be carried by the strain (Murray and Little, 1967). Resistant to *Herpes simplex virus* (Lopez, 1975). Resistant to herpes simplex virus-1 (contrast BALB/c) (Brenner *et al*, 1994). Susceptible to mouse hepatitis virus type 3 infection (Le Prevost *et al*, 1975).

Develops antibodies to mouse hepatitis virus which can be reliably detected by the complement fixation test, in contrast to five other strains (Kagiyama *et al*, 1991). Low mortality in a natural epizootic of ectromelia (Briody, 1966). High expression of RNA tumor virus group-specific antigen in some substrains but low in others (Whitmire and Salerno, 1972). Resistant to development of leukemia on infection by Friend virus (Dietz and Rich, 1972). Resistant to

diabetogenic effects of encephalomyocarditis virus, but treatment with carrageenan to compromise macrophage function makes the mice susceptible (Hirasawa *et al*, 1995). Susceptible to measles virus induced encephalitis, which correlates with a high cytotoxic T-lymphocyte response (like C3H, contrast BALB/c) (Niewiesk *et al*, 1993). Resistant to the development of tumors following inoculation with polyoma virus, in contrast with C3H/Bi (Freund *et al*, 1992). Resistant to the development of chronic Chagas' cardiomyopathy in postacute *Trypanosoma cruzi* infection (Rowland *et al* 1992). Resistant to infection with *Trypanosoma congolense* with an initial peak of parasitemia on day six, followed by rapid apparent clearance in an average of three days (contrast BALB/c) (Ogunremi and Tabel, 1995). Infection with larval *Echinococcus multilocularis* by transportal injection of hyatid homogenate results in a multivesiculation form of hyatid development (Nakaya *et al*, 1997). Susceptible to mouse adenovirus type 1 which causes a fatal hemorrhagic encephalomyelitis (contrast BALB/c) (Guida *et al*, 1995). Less susceptible to *Streptococcus suis* type 2 including the type strain, two isolates from meningitis in pigs and two isolates from tonsils of clinically healthy pigs (Kataoka *et al*, 1991). Resistant to carditis on infection with Lyme borreliosis (*Borrelia burgdorferi*) (contrast C3H, SWR, BALB/c) (Barthold *et al*, 1990). Thymectomized C57BL/6 mice that were intravenously infused with monoclonal antibody to selectively deplete CD4+ T cells are susceptible to disseminated *Mycobacterium avium* infection. The increased susceptibility is comparable to that of C57BL/6-bg. The course of such infections can be markedly restrained and in some cases the infections can be sterilized by treatment over a 120-day period with the antimycobacterial agent rifabutin (Furney *et al*, 1990). Susceptible to infection with *M. avium* strains 101 and 2-151, and can be used to test anti-mycobacterial agents (Furney *et al*, 1995). Susceptible to infection with *M. paratuberculosis* (contrast C3H/HeJ) (Tanaka *et al*, 1994). Resistant to infection with *Yersinia enterocolitica* associate with a good interferon gamma response (contrast BALB/c) (Autenrieth *et al*, 1994). Susceptible, with high amylase response to the fungus *Paracoccidioides brasiliensis* (Xidieh *et al*, 1994). Mouse mammary tumor proviral loci have been identified by Lee and Eicher (1990). Resistant to infection with *Ehrlichia risticii* (Williams and Timoney, 1994). Highly susceptible to *Plasmodium berghei* with all mice developing erythrocytic infection following intravenous injection of 50 sporozoites. The same level of infection could only be established in BALB/c with 10,000 sporozoites (Scheller *et al*, 1994). Infection with *P. berghei* results in low blood parasitemia and death with neurological symptoms within eight-ten days, in contrast with the more resistant BALB/c (Moumaris *et al*, 1995).

Resistant to chronic weakness and inflammation following infection with Tucon strain of coxsackie virus B1, in contrast with C57BL/10 and B10 congenic strains (Tam and Messner, 1996).

Life-span and spontaneous disease

Primary lung tumors 1% in males, 3% in breeding females and zero in virgin females. Lymphatic leukemia less than 2%, mammary adenocarcinomas less than 1% (Hoag, 1963). Leukemia 7% (Myers *et al*, 1970). Rare "lipomatous" hamartomas or choristomas have been noted (Adkison *et al*, 1991). Susceptible to the development of atheromatous lesions on wall of aorta after 20 weeks on a high-fat diet (Thompson, 1968; Roberts and Thompson, 1976). Develop fatty streak-like lesions in the valve sinus region of the ascending aorta after 10-20 weeks on a diet enriched in saturated fat and cholesterol. After a further 15 weeks fibro-fatty lesions with many of the characteristics of human atheromatous plaques are found (Stewart-Phillips and Lough 1991).

Exhibit aortic cartilaginous metaplasia (contrast C3H) (Qiao *et al*, 1995). Susceptible to diet-induced aortic fatty streak lesions which correlates with a low level of paroxinase mRNA (contrast C3H) (Shih *et al*, 1996). Develops non-insulin-dependent diabetes mellitus and hypertension when fed a high fat-high simple carbohydrate diet, whereas A/J mice do not (Mills *et al*, 1993). Susceptible to the development of atherosclerosis on a semi-synthetic high fat diet (Nishina *et al*, 1993). Blood glucose levels and insulin insensitivity in crosses between diet-induced type II diabetes sensitive C57BL/6 and resistant A/J are genetically independent (Surwit *et al*, 1991). High simple carbohydrate diet for five months induced hyperglycemia, hyperinsulinemia and hypercholesterolemia and non-insulin-dependent diabetes mellitus which appeared to be associated with the metabolic characteristics of visceral fat (Rebuffe-Scrive *et al*, 1993). Gain more weight on high fat diets without consuming more calories than A/J mice and develop adipocyte hyperplasia. However, animals fed a low fat, high sucrose diet were leaner than those fed a high-complex-carbohydrate diet. These results suggest that genetic differences in metabolic response to fat are more important in the development of obesity and diabetes than caloric intake (Surwit *et al*, 1995). Loci on chromosomes 1, 3, 5 and 11 are associated with variation in high density lipoprotein levels with coordinate expression of cholesterol-7-alpha hydroxylase in a cross involving atherosclerosis resistant C3H/HeJ mice (Machleder *et al*, 1997). Hepatic stearoyl CoA desaturase mRNA levels significantly elevated compared with atherosclerosis-resistant BALB/c mice, and was reduced in mice fed a high fat diet (Park *et al*, 1997). Congenital abnormalities 10%, including eye defects, polydactyly and otocephaly (Kalter, 1968). Microphthalmia and anophthalmia 8-20% and hydrocephalus 1-3% (Dagg, 1966). Ocular defects appear to be due to defects in development of the lens (Robinson *et al*, 1993). Develop spontaneous auditory degeneration with onset during young adulthood, with enhanced susceptibility to acoustic injury and delayed effects of toluene (contrast CBA/Ca) (Li, 1992, Willott *et al*, 1993; Li *et al*, 1993; Li and Borg, 1993). This is associated with early hair cell changes including bent and fused stereocilia, bulging of the cuticle plates, hair cell loss and swelling of affected dendrites (Hultcrantz and Li, 1993).

C57BL/6 mice carry a single recessive gene different from that found in BALB/cBy and WB/ReJ, causing age-related hearing loss (Willott *et al*, 1995). Hearing loss is caused by degeneration of the organ of Corti, originating in the basal, high frequency region and then proceeding apically over time. This results in a severe sensorineural hearing loss by 14 months of age (Walton *et al*, 1995). More susceptible to noise-induced hearing loss than CBA/J (Erway *et al*, 1996).

Median life-span 22.4 months in C57BL/6 males and 23.6 months in C57BL/6 females (Storer, 1966). Median life-span 24.7 and 29.6 months in C57BL/6 males and 23.6 and 29.8 months in C57BL/6 females (Les, 1969). Median life-span 27.6 months in C57BL/6 males and 27.3 months in C57BL/6 females (Goodrick, 1975). Median life-span 29.3 months in C57BL/6 males and 26.5 months in C57BL/6 females (Kunstyr and Leuenberger, 1975). Median life-span 20.0 months (Curtis, 1971). Gross tumor incidence 70%, maximum life-span about 40 months in SPF conditions (Mewissen, 1971).

Dermatitis with intense pruritis leading to self-mutilation and death, and sometimes associated with the mite *Myobia musculi* appears to be more severe in this strain than others (Csiza and McMartin, 1976). Impaired axonal regeneration involving multiple genetic loci (Lu *et al*, 1994)

Miscellaneous

High degree of genetic distinctiveness (Taylor, 1972). Recommended host for the following transplantable tumors: mammary adenocarcinoma BW 10232 melanoma B16, myeloid leukemia C 1498 and preputial gland carcinoma ESR586 (Kaliss, 1972). Embryonic stem cell lines have been established (Kawase *et al*, 1994). High rate of spontaneous mutations at the agouti and W loci (Schlager and Dickie, 1967). Characteristics of the A strain have been described by Festing (1997) and Lyon *et al*, (1996).

Physiology and biochemistry

Low plasma cholesterol at 12 and 24 weeks (Weibust, 1973). Low plasma triglyceride levels (in By and in J substrains) and low plasma cholesterol (in By and in J substrains) (Jiao *et al*, 1990). Low serum ceruloplasmin levels in males but intermediate in females (Meier and MacPike, 1968). High blood sugar (Nishimura, 1969). Low serum cholesterol in C57BL/6-a'a (Bruell *et al*, 1962). Arterial blood has a low pH (Bernstein, 1966). Low concentration of prostaglandin F in epididymis (Badr, 1975). High liver tyrosine aminotransferase in fasted mice but low in C57BL/6-ob (Blake, 1970). Low brain γ -glutamic acid decarboxylase (GAD) and

acetylcholinesterase activity but high catechol-O-methyltransferase activity (Tunncliff *et al*, 1973). Low calcium uptake by the heart (Mokler and Iturrian, 1973). Low sensitivity to thyrotropin (Levy *et al*, 1965). High brain sulphatide (Sampugna *et al*, 1975). High hepatic benz (alpha) pyrene hydroxylase activity (Kodama and Bock, 1970). Low hepatic delta-aminolaevulinic acid dehydratase activity (Doyle and Schimke, 1969). High aldehyde dehydrogenase and alcohol dehydrogenase activity compared with DBA/2 (Sheppard *et al*, 1968). High metabolism of ^{131}I with low 48 h retention (Chai *et al*, 1957). High liver arylsulphatase activity (Daniel, 1976). Low porphyrin content of Harderian gland (Margolis, 1971). Low hepatic urokinase activity but high hepatic histidine ammonia-lyase activity (Hanford *et al*, 1974).

Low basal levels of kidney catalase, superoxide dismutase and renal glutathione reductase (Misra *et al*, 1991). Iron overload causes inhibition of hepatic uroporphyrinogen decarboxylase and uroporphyrin in C57BL/10ScSn but not DBA/2 mice. This was not correlated with the Ah locus in a study involving 12 mouse strains (Smith and Francis, 1993). Low levels of apoA-IV messenger RNA in liver compared with 129/J (Reue *et al*, 1993). Low susceptibility to audiogenic seizures (Deckard *et al*, 1976). Long tau DD, the endogenous (free-running) period of the circadian pacemaker measured in constant environmental darkness (Schwartz and Zimmerman 1990). Has defective secretory group II phospholipase A2 gene (cf strains 129/Sv and B10. RIII) (Kennedy *et al*, 1995). Susceptible to severe hypercapnia with hypoxia assessed by elevated minute ventilation rate (Tankersley *et al*, 1994). Has a rapid and shallow breathing pattern phenotype (contrast C3H) (Tankersley *et al*, 1997). Susceptible to cerebral ischemia following bilateral carotid occlusion with 90% of mice showing typical neurological signs such as torsion of the neck and rolling fits with selective neuronal death in the hippocampus and caudoputamen after 20 minutes of ischemia (Yang *et al*, 1997).

Reproduction

Good reproductive performance. Litter size 6.2, sterility 8% (Nagasawa *et al*, 1973). Large litter size, mean 6.2 (Verley *et al*, 1967). Good breeding performance, 2.5 young/female/ month (Hansen *et al*, 1973). Has longer and more regular estrus cycles than DBA/2 and C3H/HeJ (Nelson *et al*, 1992). Late opening of vagina and first cornification, but early onset of cyclicity compared with C3H (Nelson *et al*, 1990). The cleavage of preimplantation embryos is faster in C57BL/6 mice than in CBA/Ca mice (McElhinny *et al*, 1996).

References

- Adkison DL, Sundberg JP (1991) "Lipomatous" hamartomas and choristomas in inbred laboratory mice. *Vet. Pathol.* 28, 305-312.
- Akamatsu Y, Barton (1974) Neoplasm and amyloidosis in strains of mice treated with 3-methylcholanthrene. *J. Nat. Cancer Inst.* 52, 377
- Almeida RMA, Correa B, Xavier JG, Mallozzi MAB, Gambale W, Paula CR (1996) Acute effect of aflatoxin B-1 on different inbred mouse strains. *Mycopathologia* 133, 23-29.
- Ammassari-Teule M, Hoffman HJ, Rossi-Arnaud C (1993) Learning in inbred mice: strain-specific abilities across three radial maze problems. *Behav. Genet.* 23, 405-412.
- Anderson LM, Jones AB, Kovatch RM (1990) Effect of pretreatment with beta-naphthoflavone on tumorigenesis by N-nitrosoethylurea in five mouse strains. *Cancer Lett.* 16, 91-94.
- Autenrieth IB, Beer M, Bohn E, Kaufmann SHE, Heesemann J (1994) Immune responses to *Yersinia enterocolitica* in susceptible BALB/c and resistant C57BL/6 mice: An essential role for gamma interferon. *Infect. Immun.* 62, 2590-2599.
- Babbini M, Pong SF, King WT, White CL (1974) Mobility of mice after amphetamine: effects of strain aggregation and illumination. *Pharmacol. Biochem. Behav.* 2, 803-809.
- Badr FM (1975) Prostaglandin levels in tissues of the male reproductive system in six strains of mice. *Endocrinol.* 96, 540-543.
- Barthold SW, Beck DS, Hansen GM, Terwilliger GA, Moody KD (1990) Lyme borreliosis in selected strains and ages of laboratory mice. *J. Infect. Dis.* 162, 133-138.
- Beamer WG, Donahue LR, Rosen CJ, Baylink DJ (1996) Genetic-variability in adult bone-density among inbred strains of mice. *Bone* 18, 397-403.
- Beamer WG, Schultz KL, Donahue LR, Churchill GA, Sen S, Wergedal JR, Baylink DJ, Rosen CJ (2001) Quantitative trait loci for femoral and lumbar vertebral bone mineral density in C57BL/6J and C3HeJ inbred strains of mice. *J. Bone Miner. Res.* 16, 1195-1206.
- Beauchamp GK, Fisher AS (1993) Strain differences in consumption of saline solutions by mice. *Physiol. Behav.* 54, 179-184.
- Belknap JK, Crabbe JC, Riggan J, O'Toole LA (1993) Voluntary consumption of morphine in 15 inbred mouse strains. *Psychopharmacology* 112, 352-358.
- Bennett LM, Farnham PJ, Drinkwater NR (1995) Strain-dependent differences in DNA synthesis and gene expression in the regenerating livers of C57BL/6J and C3H/HeJ mice. *Molecular Carcinogenesis* 14, 46-52.
- Bernstein SE (1966) Physiological characteristics. In: *Biology of the Laboratory Mouse*. 2nd. ed. (Green EL, ed). New York: McGraw-Hill, pp. 337-350.
- Bhathal PS, Jordan TW, Mackay IR (1990) Mouse strain differences in susceptibility to sporidemin-induced biliary tract injury. *Liver* 10, 193-204.
- Blake RL (1970) Regulation of liver tyrosine amino transferase activity in inbred strains and mutant mice. I. Strain variance in fasting enzyme levels. *Int. J. Biochem.* 1, 361-370.
- Blomberg B, Geckeler WR, Weigert M (1972) Genetics of the antibody response to Dextran in mice. *Science* 177, 178-180.
- Bovet D, Bovet-Nitti F, Oliverio A (1966) Effects of nicotine on avoidance conditioning of inbred strains of mice. *Psychopharmacologia* 10, 1-5.
- Bralley HC, Freeman MJ (1971) Strain differences in antibody plaque-forming cell responses in inbred mice to pneumococcal polysaccharide. *Cell. Immunol.* 2, 73-81.
- Brenner GJ, Cohen N, Moynihan JA (1994) Similar immune response to nonlethal infection with herpes simplex virus-1 in sensitive (BALB/c) and resistant (C57BL/6) strains of mice. *Cell. Immunol.* 157, 510-524.
- Briody BA (1966) The natural history of mouse pox. *National Cancer Institute Monograph* 20, 105-116.
- Brooke MS (1965) Natural hemagglutinins in mice: their occurrence and properties. *Immunol.* 8, 375-383.
- Bruell JH, Daroczy AF, Hellerstein HK (1962) Strain and sex differences in serum cholesterol levels in mice. *Science* 135, 1071-1072.
- Cabib S, Bonaventura N (1997) Parallel strain-dependent susceptibility to environmentally-induced stereotypes and stress-induced behavioral sensitisation in mice. *Physiol. Behav.* 61, 499-506.
- Cartner SC, Simecka JW, Briles DE, Cassell GH, Lindsey JR (1996) Resistance to Mycoplasma lung-disease in mice is a complex genetic trait. *Infect. Immun.* 64, 5326-5331.
- Caspi RR, Ferguson TA (2012) The *rdb* mutation of the *Crb1* gene is present in vendor lines of C57BL/6N mice and embryonic stem cells, and confounds ocular induced mutant phenotypes. *IOVS Paper in Press.*
- Cerny J, McAlack RF, Sajid MA, Friedman H (1971) Genetic differences in the immunocyte response of mice to separate determinants on one bacterial antigen. *Nature New Biol.* 230, 247-248.
- Chai CK, Amin A, Reineke EP (1957) Thyroidal iodine metabolism in inbred and F1 hybrid mice. *Am. J. Physiol.* 188, 499-502.
- Chen PE, Specht CG, Morris RGM, Schoepfer R (2002) Spatial learning is unimpaired in mice containing a deletion of the alpha-synuclein locus. *European Journal of Neuroscience* 16, 154-158.
- Csiza CK, McMartin DN (1976) Apparent acral dermatitis in a C57BL/6 Nya mouse colony. *Lab. Animal Sci.* 26, 781-787.
- Curtis HJ (1971) Genetic factors in aging. *Adv. Genet.* 16, 305-324.
- Dagg CP (1966) Teratogenesis, in *Biology of the laboratory mouse*, 2nd. ed. (Green EL, ed), pp. 309-328. McGraw-Hill, New York.
- Dains K, Hitzemann B, Hitzemann R (1996) Genetics, neuroleptic response and the organization of cholinergic neurons in the mouse striatum. *J. Pharmacol. Exp. Therapeut.* 279, 1430-1438.
- Daniel WL (1976) Genetics of murine liver and kidney arylsulfatase B. *Genetics* 82, 477-491.
- Davis WM, King WT (1967) Pharmacogenetic factor in the convulsive responses of mice to flurothyl. *Experientia* 23, 214-215.
- Davis WM, King WT, Rabbini M (1967) Placebo effect of saline on locomotor activity in several strains of mice. *J. Pharmacol. Sci.* 56, 1347-1349.
- Deckard BS, Lief B, Schlesinger K, DeFries JC (1976) Developmental patterns of seizure susceptibility in inbred strains of mice. *Devel. Psychobiol.* 9, 17.
- De Macario EC, Macario AJL (1980) Immunosuppression associated with erythropoiesis in genetic low responder mice. *Ann. Immunol. (Inst. Pasteur)* 131C, 397-404.
- Deringer MK, Dunn TB, Heston WE (1953) Results of exposure of strain C3H mice to chloroform. *Proc. Soc. Exp. Biol. Med.* 83, 474-479.
- Dietz M, Rick MA (1972) Effect of host strain and H-2 type on spontaneous regression of murine leukemia virus. *Int. J. Cancer* 10, 99-104.
- Di Pauli R (1972) Genetics of the immune response. I. Differences in the specificity of antibodies to lipopolysaccharides among different strains of mice. *J. Immunol.* 109, 394-400.
- Dorf ME, Dunham EK, Johnson JP, Benacerraf B (1974) Genetic control of the immune response: the effect of non-H-2 linked genes on antibody production. *J. Immunol.* 112, 1329-1336.
- Dostal M, Jelinek R (1973) Sensitivity of embryos and intraspecies differences in mice in response to prenatal administration of corticoids. *Teratology* 8, 245-252.
- Doyle D, Schimke RT (1969) The genetic and developmental regulation of hepatic L-aminolevulinatase dehydratase in mice. *J. Biol. Chem.* 244, 5449-5459.
- Erway LC, Shiau YW, Davis RR, Krieg EF (1996) Genetics of age-related hearing-loss in mice. 3. Susceptibility of inbred and F1-hybrid strains to noise-induced hearing-loss. *Hearing Research* 93, 181-187.
- Evans JG, Collins MA, Lake BG, Butler WH (1992) The histology and development of hepatic nodules and carcinoma in C3H/He and C57BL/6 mice following chronic phenobarbitone administration. *Toxicologic Pathology* 20, 585-594.
- Evans JT, Shows TB, Sproul EE, Paolini NS, Mittelman A, Hauschka TS (1977) Genetics of colon carcinogenesis in mice treated with 1, 2-dimethylhydrazine. *Cancer Res.* 37, 134-136.
- Fahey JL (1965) Differences in the electrophoretic mobility of antibody from inbred strains of mice. *J. Immunol.* 94, 819-823.
- Fawcington E, Festing MFW Mouse strain differences in balsa wood gnawing. Unpublished data.
- Festing MFW (1997) Inbred Strains of mice. *Mouse genome* 95, 519-686.
- Festing MFW, Blackmore DK (1971) Life span of specified-pathogen-free (MRC category 4) mice and rats. *Lab. Anim.* 5, 179-192.
- Finn DA, Roberts AJ, Lotrich F, Gallaher EJ (1997) Genetic differences in behavioral sensitivity to a neuroactive steroid. *J. Pharmacol. Exp. Therapeut.* 280, 820-828.
- Fox JG, Li X, Cahill RJ, Andrutis K, Rustig AK, Odze R, Wang TC (1996) Hypertrophic gastropathy in *Helicobacter felis*-infected wild-type C57BL/6 mice and p53 hemizygous transgenic mice. *Gastroenterology* 110, 155-166.
- Freund R, Dubensky T, Bronson R, Sotnikov A, Carroll J, Benjamin T (1992) Polyoma tumorigenesis in mice: Evidence for dominant resistance and dominant susceptibility genes of the host. *Virology* 191, 724-731.
- Fuchs S, Mozes E, Maoz A, Sela M (1974) Thymus independence of a collagen-like synthetic polypeptide and of collagen and the need for thymus and bone marrow-cell cooperation in the immune response to gelatin. *J. Exp. Med.* 139, 148-158.
- Fuller JL (1964) Measurement of alcohol preference in genetic experiments. *J. Comp. Physiol. Psychol.* 57, 85-88.
- Fuller JL, Sjursen FH (1967) Audiogenic seizures in eleven mouse strains. *J. Hered.* 58, 135-140.
- Furney SK, Roberts AD, Orme IM (1990) Effect of rifabutin on disseminated Mycobacterium avium infections in thymectomized, CD4 T-cell-deficient mice. *Antimicrobial Agents & Chemotherapy* 34, 1629-1632.
- Furney SK, Skinner PS, Farrer J, Orme IM (1995) Activities of rifabutin, clarithromycin and ethambutol against two virulent strains of Mycobacterium avium in a mouse model. *Antimicrobial Agents & Chemotherapy* 39, 786-789.
- Gabridge MG, Abrams GD, Murphy WH (1972) Lethal toxicity of Mycoplasma fermentans in mice. *J. Infect. Dis.* 125, 153-160.
- Gannon KS, Contreras RJ (1995) Sodium intake linked to amiloride-sensitive gustatory transduction in C57BL/6J and 129/J mice. *Physiol. Behav.* 57, 231-239.
- Gasser DL, Silvers WK (1971) Genetic basis of male skin rejection in mice. *Transplant.* 12, 412-414.
- Goldstein BD, Lai LY, Ross SR, Cuzzi-Spada R (1973) Susceptibility of inbred mouse strains to ozone. *Arch. Environ. Health* 27, 412-413.
- Goodrick CL (1975) Lifespan and the inheritance of longevity of inbred mice. *J. Gerontol.* 30, 257-263.
- Green MC, Azar CA, Maren TH (1973) Strain differences in susceptibility to the teratogenic effect of acetazolamide in mice. *Teratology* 8, 143-145.
- Griem P, Scholz E, Turfeld M, Zander D, Wiesner U, Dunemann L, Gleichmann E (1997) Strain differences in tissue concentrations of mercury in inbred mice treated with mercuric chloride. *Toxicol. Appl. Pharmacol.* 144, 163-170.
- Guida JD, Fejer G, Pirofski LA, Brosnan CF (1995) Mouse adenovirus type 1 causes a fatal hemorrhagic encephalomyelitis in adult C57BL/6 but not BALB/c mice. *Journal of Virology* 69, 7674-7681.
- Hanford WC, Nep RL, Arfin SM (1974) Genetic variation in histidine ammonia-lyase activity in the mouse. *Biochem. Biophys. Res. Comm.* 61, 1434-1437.
- Hansen CT, Judge FJ, Whitney RA (1973) Catalog of NIH rodents. National Institutes of Health. DHEW publication (NIH) 74-606, Bethesda.
- Harada M, Nagata M, Takeuchi M, Ohara T, Makino S, Watanabe A (1991) Age-dependent difference in susceptibility to IgE antibody- and IgG1 antibody-mediated passive anaphylactic shock in the mouse. *Immunological Investigations* 20, 515-523.
- Hellman A, Fowler AK (1972) Studies of the blastogenic response of murine lymphocyte. III. Specific viral transformation. *Proc. Soc. Exp. Biol. Med.* 141, 106-109.
- Hill GB, Osterhout S, O'Fallon WM (1968) Variation in response to hyperbaric oxygen among inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 129, 687-689.
- Hill RN, Clemens TL, Liu DK, Vesell ES (1975) Genetic control of chloroform toxicity in mice. *Science* 190, 159-161.
- Hirasawa K, Ogiso Y, Takeda M, Lee MJ, Itagaki S, Doi K (1995) Protective effects of macrophage-derived interferon against encephalomyelitis virus-induced diabetes mellitus in mice. *Lab. Animal Sci.* 45, 652-656.
- Hoag WG (1963) Spontaneous cancer in mice. *Ann. NY Acad. Sci.* 108, 805-831.
- Hultcrantz M, Li HS (1993) Inner ear morphology in CBA/Ca and C57BL/6J mice in relationship to noise, age and phenotype. *European Archives of Oto-Rhino-Laryngology* 250, 257-264.
- Hummel KP, Richardson FL, Fekete E (1966) *Anatomy, in Biology of the Laboratory Mouse*, 2nd. ed. (Green EL, ed), pp. 247-307. McGraw-Hill, New York.
- Iizawa Y, Wagner RD, Czuprynski CJ (1993) Analysis of cytokine mRNA expression in Listeria-resistant C57BL/6 and Listeria-susceptible A/J mice during Listeria monocytogenes infection. *Infect. Immun.* 61, 3739-3744.
- Ishii AI, Sano M (1989) Strain-dependent differences in susceptibility of mice to experimental Angiostrongylus costaricensis infection. *J. Helminthology* 63, 302-306.
- Jiao S, Cole TG, Kitchens R, Pflieger B, Schonfeld G (1990) Genetic heterogeneity of lipoproteins in inbred strains of mice: analysis by gel-permeation chromatography. *Metabolism* 39, 155-160.
- Kagiya M, Takakura A, Koyoma K, Terada E, Sakuria Y (1991) Detection of mouse hepatitis virus antibody by protein A-ELISA in 6 prevalent inbred strains or outbred stocks of mice. *Lab. Anim.* 25, 106-109.
- Kalcheva ID, Matsuda Y, Plass C, Chapman VM (1995) Isolation and characterization of a pseudoautosomal region-specific genetic-marker in C57BL/6 mice using genomic representational difference analysis. *Proceedings of the National Academy of Sciences of the United States of America* 92, 12352-12356.
- Kaliss N (1972) Transplanted tumors. *Jax Notes* 410, Jackson Laboratory, Bar Harbor, Maine.

85. Kalter H (1968) Sporadic congenital malformations of newborn inbred mice. *Teratology* 1, 193-200.
86. Kalter H (1981) Dose response studies with genetically homogeneous lines of mice as a teratology testing and risk assessment procedure. *Teratology* 24, 79-86.
87. Kataoka Y, Haritani M, Mori M, Kishima M, Sugimoto C, Nakazawa M, Yamamoto K (1991) Experimental infections of mice and pigs with *Streptococcus suis* type 2. *J. Vet. Med. Sci.* 53, 1043-1049.
88. Kawase E, Suemori H, Takahashi N, Okazaki K, Hashimoto K, Nakatsuji N (1994) Strain difference in establishment of mouse embryonic stem (ES) cell lines. *International Journal of Developmental Biology* 38, 385-390.
89. Kaye M, Kusy P (1995) Genetic lineage, bone mass and physical activity in mice. *Bone* 17, 131-135.
90. Keetch DW, Humphrey P, Ratliff TL (1994) Development of a mouse model for nonbacterial prostatitis. *Journal of Urology* 152, 247-250.
91. Kennedy BP, Payette P, Mudgett J, Vadas P, Pruzanski W, Kwan M, Tang C, Rancourt DE, Cromlish WA (1995) A natural disruption of the secretory group II phospholipase A2 gene in inbred mouse strains. *J. Biol. Chem.* 270, 22378-22385.
92. Kimura M, Takahashi H, Ohtake T, Sato T, Hishida A, Nishimura M, Honda N (1993) Interstrain differences in murine daunomycin-induced nephrosis. *Nephron* 63, 193-198.
93. Kleeberger SR, Levitt RC, Zhang LY (1993) Susceptibility to ozone-induced inflammation. I. Genetic control of the response to subacute exposure. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 264, L15-L20.
94. Kodama Y, Bock FG (1970) Benz [a] pyrene-metabolizing enzyme activity of livers of various strains of mice. *Cancer Res.* 30, 1846-1849.
95. Konno S, Adachi M, Matsuura T, Sunouchi K, Hoshino H, Okazawa A, Kobayashi H, Takahashi T (1993) Bronchial reactivity to methacholine and serotonin in six inbred mouse strains. [Japanese]. *Japanese Journal of Allergy* 42, 42-47.
96. Kosobud AE, Cross SJ, Crabbe JC (1992) Neural sensitivity to pentylentetrazol convulsions in inbred and selectively bred mice. *Brain Res.* 592, 122-128.
97. Kouri RE, Salerno RA, Whitmore CE (1973) Relationships between arylhydrocarbon hydroxylase inducibility and sensitivity to chemically induced subcutaneous sarcomas in various strains of mice. *J. Natl. Cancer Inst.* 50, 363-368.
98. Kolaczowska W, Chadzinska M, Sejelid R, Plytycz B (2000) Strain differences in some immune parameters can be obscured by circadian variations and laboratory routines: studies of male C57BL/6J, BALB/c and CB6F1 mice. *Lab. Anim.* 35, 91-100.
99. Kunstyr I, Leuenberger HGW (1975) Gerontological data on C57BL/6 mice. I. Sex differences in survival curves. *J. Gerontol.* 30, 157-162.
100. Laskay T, Diefenbach A, Rollinghoff M, Solbach V (1995) Early parasite containment is decisive for resistance to *Leishmania* major infection. *Eur. J. Immunol.* 25, 2220-2227.
101. Le AD, Ko J, Chow S, Quan B (1994) Alcohol consumption by C57BL/6, BALB/c and DBA/2 mice in a limited access paradigm. *Pharmacol. Biochem. Behav.* 47, 375-378.
102. Lee BK, Eicher EM (1990) Segregation patterns of endogenous mouse mammary tumor viruses in five recombinant inbred strain sets [published erratum appears in *J Virol* (1991) 65, 1666]. *J. Virol.* 64, 4568-4572.
103. Le Prevost C, Virelizier JL, Dupuy JM (1975) Immunopathology of mouse hepatitis virus type 3 infection. III. Clinical and virologic observation of a persistent viral infection. *J. Immunol.* 115, 640-643.
104. Les EP (1969) Personal communication and effect of pasteurized diets on lifespan of inbred mice. (Abstract 10). AALAS, Publication 69-72.
105. Levy RP, McGuire WL, Shaw RK, Bartsch GE (1965) Effect of species differences of mice on the bioassay of thyrotropin. *Endocrinol.* 76, 890-894.
106. Li HS (1992) Genetic influences on susceptibility of the auditory system to aging and environmental factors. *Scandinavian Audiology, Supplement* 21, 1-39.
107. Li HS, Borg E (1993) Auditory degeneration after acoustic trauma in two genotypes of mice. *Hearing Research* 68, 19-27.
108. Li HS, Hultcrantz M, Borg E (1993) Influence of age on noise-induced permanent threshold shifts in CBS/Ca and C57BL/6J mice. *Audiology* 32, 195-204.
109. Lopez C (1975) Genetics of natural resistance to herpes virus infections in mice. *Nature* 258, 152-153.
110. Lu X, Skamene E, Richardson PM (1994) Studies of axonal regeneration in C57BL/6J and A/J mice. *Brain Res.* 652, 174-176.
111. Lush IM (1988) The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet. Res.* 53, 95-99.
112. Lyon MF, Rastan S, Brown SDM (1996) Genetic variants and strains of the laboratory mouse. 2 Volumes. Oxford, New York, Tokyo: Oxford University Press.
113. Macario AJL, Stahl W, Miller RM (1980) Lymphocyte subpopulations and function in chronic murine toxoplasmosis. II. Cyclic immunosuppression in genetic-low-responder mice. *Cell. Immunol.* 56, 235-239.
114. Machleder D, Ivandic B, Welch C, Castellani L, Reue K, Lusa AJ (1997) Complex genetic control of HDL levels in mice in response to an atherogenic diet - Coordinate regulation of HDL levels and bile acid metabolism. *J. Clin. Invest.* 99, 1406-1419.
115. Margolis FL (1971) Regulation of porphyrin biosynthesis in the Harderian gland of inbred mouse strains. *Arch. Biochem. Biophys.* 145, 373-381.
116. Marks MJ, Stitzel JA, Collins AC (1989) Genetic influences on nicotine responses. *Pharmacol. Biochem. Behav.* 33, 667-678.
117. McCarthy MM, Dutton RW (1975) The humoral response of mouse spleen cells to two types of sheep erythrocytes. *J. Immunol.* 115, 1316-1321.
118. McClive PJ, Huang D, Morahan G (1994) C57BL/6 and C57BL/10 inbred mouse strains differ at multiple loci on chromosome 4. *Immunogenet.* 39, 286-288.
119. McElhinny AS, Davis FC, Warner CM (1996) The effect of melatonin on cleavage rate of C57BL/6 and CBA/Ca preimplantation embryos cultured in vitro. *J. Pineal Res.* 21, 44-48.
120. McMillan BA, Williams HL (1998) Role of taste and calories in the selection of ethanol by C57BL/6N-Hsd and Hsd:ICR mice. *Alcohol* 15, 193-198.
121. Meier H, MacPike AD (1968) Levels and heritability of serum ceruloplasmin activity in inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 128, 1185-1190.
122. Melo JA, Shendure J, Pociask K, Silver LM (1996) Identification of sex-specific quantitative trait loci controlling alcohol preference in C57BL/6 mice. *Nature Genet.* 13, 147-153.
123. Mendoza LA, Hamburg M, Fuld H (1967) Differences in thyroid activity in several inbred strains of mice. *Anat. Rec.* 158, 275-280.
124. Messeri P, Oliverio A, Bovet D (1972) Relations between avoidance and activity. A diallel study in mice. *Behav. Biol.* 7, 733-742.
125. Mewissen DJ (1971) Natural tumor incidence in a population of mice as a reference index. *Fed. Proc.* 30, 311.
126. Militzer K, Wecker E (1986) Behavior-associated alopecia areata in mice. *Lab. Anim.* 20, 9-13.
127. Mills E, Kuhn CM, Feinglos MN, Surwit R (1993) Hypertension in CB57BL/6J mouse model of non-insulin-dependent diabetes mellitus. *Am. J. Physiol.* 264, R73-R78.
128. Misra M, Rodriguez RE, North SL, Kasprzak KS (1991) Nickel-induced renal lipid peroxidation in different strains of mice: concurrence with nickel effect on antioxidant defence systems [published erratum appears in *Toxicol. Lett.* (1992) 60, 239]. *Toxicol. Lett.* 58, 121-133.
129. Miyake T, Cameron AM, Hall BK (1996a) Detailed staging of inbred C57BL/6 mice between Theiler's [1972] stages 18 and 21 (11-13 days of gestation) based on craniofacial development. *Journal of Craniofacial Genetics and Developmental Biology* 16, 1-31.
130. Miyake T, Cameron AM, Hall BK (1996b) Stage-specific onset of condensation and matrix deposition for Meckel's and other first arch cartilages in inbred C57BL/6 mice. *Journal of Craniofacial Genetics and Developmental Biology* 16, 32-47.
131. Mohammadi M, Redline R, Nedrud J, Cizin S (1996) Role of the host in pathogenesis of *Helicobacter*-associated gastritis: *H. felis* infection of inbred and congenic mouse strains. *Infect. Immun.* 64, 238-245.
132. Mokler CM, Iturrian WB (1973) Strain differences in subcellular calcium distribution in striated muscle of the mouse. *Proc. Soc. Exp. Biol. Med.* 142, 919-923.
133. Monroy-Ostria A, Fuentes-Fraga I, Garcia-Flores C, Favila-Castillo L (1994) Infection of BALB/c, C57BL/6 mice and F1 hybrid CB6F1 mice with strains of *Leishmania mexicana* isolated from Mexican patients with localized or diffuse cutaneous leishmaniasis. *Archives of Medical Research* 25, 401-406.
134. Moumaris M, Sestier C, Miltgen F, Halbreich A, Gentilini M, Sabolovic D (1995) Effect of fatty acid treatment in cerebral malaria-susceptible and nonsusceptible strains of mice. *Journal of Parasitology* 81, 997-999.
135. Muller-Sieburg CE, Riblet R (1996) Genetic control of the frequency of hematopoietic stem cells in mice: Mapping of a candidate locus to chromosome 1. *Journal Of Experimental Medicine* 183, 1141-1150.
136. Murray WS, Little CC (1967) Genetic studies of carcinogenesis in mice. *J. Natl. Cancer Inst.* 38, 639-656.
137. Muthing J (1997) Neutral glycosphingolipids and gangliosides from spleen T lymphoblasts of genetically different inbred mouse strains. *Glycoconjugate Journal* 14, 241-248.
138. Myers DD, Meier H, Huebner RJ (1970) Prevalence of murine C-type RNA virus group specific antigen in inbred strains of mice. *Life Sci.* 9, 1071-1080.
139. Nagasawa H, Miyamoto M, Fujimoto M (1973) Reproductivity in inbred strains of mice and project for their efficient production. *Exp. Animals (Japan)* 22, 119-126.
140. Nakaya K, Nakao M, Ito A (1997) *Echinococcus multilocularis*: Mouse strain difference in hydatid development. *J. Helminthology* 71, 53-56.
141. Nelson JF, Karelus K, Felicio LS, Johnson TE (1990) Genetic influences on the timing of puberty in mice. *Biol. Reprod.* 42, 649-655.
142. Nelson JF, Karelus K, Felicio LS, Johnson TE (1992) Genetic influences on estrous cyclicity in mice: evidence that cycle length and frequency are differentially regulated. *J. Reprod. Fertil.* 94, 261-268.
143. Neuhaus IM, Sommarhald CS, Johnson DK, Beier DR (1997) Microsatellite DNA variants between the FVB/N and C3HeB/FeJLa and C57BL/6J mouse strains. *Mammalian Genome* 8, 506-509.
144. Ng GYK, Odowd BF, George SR (1996) Genotypic differences in mesolimbic enkephalin gene-expression in DBA/2J and C57BL/6J inbred mice. *Eur. J. Pharmacol.* 311, 45-52.
145. Niewiesk S, Brinckmann U, Bankamp B, Sirak S, Liebert UG, Ter Meulen V (1993) Susceptibility to measles virus-induced encephalitis in mice correlates with impaired antigen presentation to cytotoxic T lymphocytes. *Journal of Virology* 67, 75-81.
146. Nikulina EM, Skrinskaya JA, Popova N. K (1991) Role of genotype and dopamine receptors in behavior of inbred mice in a forced swimming test. *Psychopharmacology* 105, 525-529.
147. Nishi Y, Hasegawa MM, Inui N (1993) Genetic variations in baseline and ultraviolet light-induced sister chromatid exchanges in peritoneal lymphocytes among different mouse strains. *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis* 286, 145-154.
148. Nishimura M (1969) Breeding of mice strains for diabetes mellitus. *Exp. Animals (Japan)* 18, 147-157.
149. Nishina PM, Wang J, Toyofuku W, Kuypers FA, Ishida BY, Paigen B (1993) Atherosclerosis and plasma and liver lipids in nine inbred strains of mice. *Lipids* 28, 599-605.
150. Nishioka Y. (1987) Y-chromosomal DNA polymorphism in mouse inbred strains. *Genet. Res.* 50, 69-72.
151. Noonan FP, Hoffman HA (1994) Susceptibility to immunosuppression by ultraviolet B radiation in the mouse. *Immunogenet.* 39, 29-39.
152. Oesch F, Morris N, Daly JW (1973) Genetic expression of the induction of epoxide hydrolase and aryl hydrocarbon hydroxylase activities in the mouse by phenobarbital or 3-methylcholanthrene. *Molec. Pharmacol.* 9, 692-696.
153. Ogunremi O, Tabel H (1995) Genetics of resistance to *Trypanosoma congolense* in inbred mice: Efficiency of apparent clearance of parasites correlates with long-term survival. *Journal of Parasitology* 81, 876-881.
154. Otterness DM, Weinsilboum RM (1987a) Mouse thiopurine methyltransferase pharmacogenetics: biochemical studies and recombinant inbred strains. *J. Pharmacol. Exp. Therapeut.* 240, 180-186.
155. Otterness DM, Weinsilboum RM (1987b) Mouse thiopurine methyltransferase pharmacogenetics: Monogenic inheritance. *J. Pharmacol. Exp. Therapeut.* 240, 817-824.
156. Paquette NC, Zhang LY, Ellis WA, Scott AL, Kleeberger SR (1996) Vitamin A deficiency enhances ozone-induced lung injury. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 270, L475-L482.
157. Park EI, Paisley EA, Mangian HJ, Swartz DA, Wu MX, O'Morchoe PJ, Behr SR, Visek WJ, Kaput J (1997) Lipid level and type alter stearyl CoA desaturase mRNA abundance differently in mice with distinct susceptibilities to diet-influenced diseases. *J. Nutr.* 127, 566-573.
158. Pinchuck P, Maurer PH (1965) Antigenicity of polypeptides (poly alpha amino acids). XVI. Genetic control of immunogenicity of synthetic polypeptides in mice. *J. Exp. Med.* 122, 673-679.
159. Qiao JH, Fishbein MC, Demer LL, Lusa AJ (1995) Genetic determination of cartilaginous metaplasia in mouse aorta. *Arteriosclerosis, Thrombosis and Vascular Biology* 15, 2265-2272.
160. Rebuffe-Scrive M, Surwit R, Feinglos M, Kuhn C, Rodin J (1993) Regional fat distribution and metabolism in a new mouse model (C57BL/6J) of non-insulin-dependent diabetes mellitus. *Metabolism: Clinical and Experimental* 42, 1405-1409.
161. Reue K, Purcell-Huynh DA, Leete TH, Doolittle MH, Durstenfeld A, Lusa AJ (1993) Genetic variation in mouse apolipoprotein A-IV expression is determined pre- and post-transcriptionally. *J. Lipid Res.* 34, 893-903.
162. Roberts A, Thompson JS (1976) Inbred mice and their hybrids as an animal model for atherosclerosis research. In: *Atherosclerosis Drug Discovery* (Day CE, ed). New York: Plenum Press, pp. 313-327.
163. Robinson ML, Holmgren A, Dewey MJ (1993) Genetic control of ocular morphogenesis: Defective lens development associated with ocular anomalies in C57BL/6 mice. *Exp. Eye Res.* 56, 7-16.
164. Robinson SF, Marks MJ, Collins AC (1996) Inbred mouse strains vary in oral self-selection of nicotine. *Psychopharmacology* 124, 332-339.
165. Robson HG, Vas SI (1972) Resistance of mice to *Salmonella typhimurium*. *J. Infect. Dis.* 126, 378-380.

166. Rodgers DA (1966) Factors underlying differences in alcohol preference among inbred strains of mice. *Psychosomat. Med.* 28, 498-513.
167. Rowland EC, Lozykowski MG, McCormick TS (1992) Differential cardiac histopathology in inbred mouse strains chronically infected with *Trypanosoma cruzi*. *Journal of Parasitology* 78, 1059-1066.
168. Rubinstein P, Liu N, Strenn EW, Decary F (1974) Electrophoretic mobility and agglutinability of red blood cells: a 'new' polymorphism in mice. *J. Exp. Med.* 139, 313-322.
169. Russell ES, Neufeld EF, Higgins CT (1951) Comparison of normal blood picture of young adults from 18 inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 78, 761-766.
170. Sadarangani C, Skamene E, Kongshaven PAL (1980) Cellular basis for genetically determined enhanced resistance of certain mouse strains to Listeriosis. *Infect. Immun.* 28, 381-386.
171. Sakagami T, Dixon M, O'Rourke J, Howlett R, Alderuccio F, Vella J, Shimoyama T, Lee A (1996) Atrophic gastric changes in both *Helicobacter felis* and *Helicobacter pylori* infected mice are host dependent and separate from antral gastritis. *Gut* 39, 639-648.
172. Sampugna J, Clements J, Carter TP, Campagnoni AT (1975) Comparison of lipids in total brain tissue from five mouse genotypes. *J. Neurobiol.* 6, 259-266.
173. Sato SI, Takizawa H, Inui N (1993) Mouse strain differences in induction of micronuclei by base analogues and nucleosides. *Mutation Research - Mutation Research Letters* 301, 45-49.
174. Scheller LF, Wirtz RA, Azad AF (1994) Susceptibility of different strains of mice to hepatic infection with *Plasmodium berghei*. *Infect. Immun.* 62, 4844-4847.
175. Schlager G (1968) Kidney weight in mice: strain differences and genetic determination. *J. Hered.* 59, 171-174.
176. Schlager G, Dickie MM (1967) Spontaneous mutations and mutation rates in the house mouse. *Genetics* 57, 319-330.
177. Schlesinger K, Wimer R (1967) Genotype and conditioned avoidance learning in the mouse. *J. Comp. Physiol. Psychol.* 63, 139-141.
178. Schmid FA, Dickey PA, Stanco GA, Tarnowski GS (1966) Toxicity of intraperitoneal injections of 7,12-dimethylbenz (a) anthracene (DMBA) and other agents in inbred mice. *Proc. Am. Assoc. Cancer Res.* 7, 62.
179. Schwartz WJ, Zimmerman P (1990) Circadian timekeeping in BALB/c and C57BL/6 inbred mouse strains. *Journal of Neuroscience* 11, 3685-3694.
180. Scott P, Eaton A, Gause WC, Zhou XD, Hondowicz B (1996) Early IL-4 production does not predict susceptibility to *Leishmania major*. *Experimental Parasitology* 84, 178-187.
181. Shepard CC, Habas JA (1967) Relation of infection to tissue temperature in mice infected with *Mycobacterium marinum* and *Mycobacterium leprae*. *J. Bacteriol.* 93, 790-796.
182. Sheppard JR, Albersheim P, McClean GE (1968) Enzyme activities and ethanol preference in mice. *Biochem. Genet.* 2, 205-212.
183. Shih DM, Gu L, Hama S, Xia YR, Navab M, Fogelman AM, Lusis AJ (1996) Genetic-dietary regulation of serum paraoxonase expression and its role in atherogenesis in a mouse model. *J. Clin. Invest.* 97, 1630-1639.
184. Slingsby JH, Hogarth MB, Simpson E, Walport MJ, Morley BJ (1996) New microsatellite polymorphisms identified between C57BL-6, C57BL-10 and C57BL-KsJ inbred mouse strains. *Immunogenet.* 43, 72-75.
185. Smith AG, Francis JE (1993) Genetic variation of iron-induced uroporphyrin in mice. *Biochem. J.* 291, 29-35.
186. Specht CG, Schoepfer R (2001) Deletion of the alpha-synuclein locus in a subpopulation of C57BL/6J mice. *BioMed Central Neuroscience* 2, 11.
187. Srivastava RAK (1995) Increased apoB100 mRNA in inbred strains of mice by estrogen is caused by decreased RNA editing protein mRNA. *Biochemical and Biophysical Research Communications* 212, 381-387.
188. Steinberg AD, Pincus T, Talal N (1971) The pathogenesis of autoimmunity in New Zealand mice. III. Factors influencing the formation of anti-nucleic acid antibodies. *Immunol.* 20, 523-531.
189. Stewart-Phillips JL, Lough J (1991) Pathology of atherosclerosis in cholesterol-fed, susceptible mice. *Atherosclerosis* 90, 211-218.
190. Storer JB (1966) Longevity and gross pathology at death in 22 inbred strains of mice. *J. Gerontol.* 21, 404-409.
191. Surwit RS, Seldin MF, Kuhn CM, Cochran C and Feinglos MN (1991) Control of expression of insulin resistance and hyperglycemia by different genetic factors in diabetic C57BL/6J mice. *Diabetes* 40, 82-87.
192. Surwit RS, Feinglos MN, Rodin J, Sutherland A, Petro AE, Opara EC, Kuhn CM, Rebuffe-Scrive M (1995) Differential effects of fat and sucrose on the development of obesity and diabetes in C57BL/6J and A/J mice. *Metabolism: Clinical and Experimental* 44, 645-651.
193. Takahashi M, Kleeberger SR, Croxton TL (1995) Genetic control of susceptibility to ozone-induced changes in mouse tracheal electrophysiology. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 269, L6-L10.
194. Tam PE, Messner RP (1996) Genetic-determinants of susceptibility to coxsackievirus B1-induced chronic inflammatory myopathy - effects of host background and major histocompatibility complex genes. *Journal of Laboratory and Clinical Medicine* 128, 279-289.
195. Tanaka S, Sato M, Taniguchi T, Yokomizo Y (1994) Histopathological and morphometrical comparison of granulomatous lesions in BALB/c and C3H/HeJ mice inoculated with *Mycobacterium paratuberculosis*. *J. Comp. Pathol.* 110, 381-388.
196. Tanioka Y, Esaki K (1971) Strain differences in mortality of anaphylactic shock in mice-challenging by intravenous injection. *Exp. Animals (Japan)* 20, 127-130.
197. Tankersley CG, Fitzgerald RS, Kleeberger SR (1994) Differential control of ventilation among inbred strains of mice. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology* 267, R1371-R1377.
198. Tankersley CG, Fitzgerald RS, Levitt RC, Mitzner WA, Ewart SL, Kleeberger SR (1997) Genetic control of differential baseline breathing pattern. *J. Appl. Physiol.* 82, 874-881.
199. Taylor BA (1972) Genetic relationship between inbred strains of mice. *J. Hered.* 63, 83-86.
200. Taylor BA (1976) Genetic analysis of susceptibility to isoniazid-induced seizures in mice. *Genetics* 83, 373-377.
201. Thomas PE, Hutton JJ, Taylor BA (1973) Genetic relationship between aryl hydrocarbon hydroxylase inducibility and chemical carcinogen induced skin ulceration in mice. *Genetics* 74, 655-659.
202. Thompson JS (1968) Atherosclerosis in inbred strains of mice. *Anat. Res.* 160, 440.
203. Thompson JS (1969) Atheromata in an inbred strain of mice. *J. Atheroscler. Res.* 10, 113-122.
204. Thompson WR (1953) The inheritance of behavior: behavioral differences in fifteen mouse strains. *Can. J. Psychol.* 7, 145-155.
205. Toda S, Kimura M, Tohya K (1989) Strain differences in histamine release from mouse peritoneal mast cells induced by compound 48/80 or A23187. *Jikken Dobutsu - Experimental Animals* 38, 135-137.
206. Treadwell PE (1969) The inheritance of susceptibility to anaphylaxis in inbred mice and their hybrid progenies. *J. Reticuloendothel. Soc.* 6, 343-353.
207. Tunncliffe G, Wimer CC, Wimer RE (1973) Relationships between neurotransmitter metabolism and behavior in seven inbred strains of mice. *Brain Res.* 61, 428-434.
208. Vadasz C, Fleischer A, Lafrancois J, Mao RF (1996) Self-administration of ethanol - towards the location of predisposing polygenes in quasi-congenic animal-models. *Alcohol* 13, 617-620.
209. Verley FA, Grahm D, Leslie WP, Hamilton KF (1967) Sex ratio of mice as possible indicator of mutation rate for sex-linked lethals. *J. Hered.* 58, 285-290.
210. Wahlsten D (1973) Contribution of the genes albinism (c) and retinal degeneration (rd) to a strain-by-training procedure interaction in avoidance learning. *Behav. Genet.* 3, 303-316.
211. Walton JP, Frisina RD, Meierhans LR (1995) Sensorineural hearing loss alters recovery from short-term adaptation in the C57BL/6 mouse. *Hearing Research* 88, 19-26.
212. Weibust RS (1973) Inheritance of plasma cholesterol levels in mice. *Genetics* 73, 303-312.
213. Weissman I (1967) Genetic and histochemical studies on mouse spleen black spots. *Nature* 215, 315.
214. Whitmire CE, Salerno RA (1972) RNA tumor virus antigen and tumor induction by various doses of 3-methylcholanthrene in various strains of mice treated as weanlings. *Cancer Res.* 32, 1129-1132.
215. Whitmire CE, Salerno RA, Rabstein LS, Heubner RJ, Turner HC (1971) RNA tumor-virus antigen expression in chemically induced tumors. Virus-genome specified common antigens detected by complement fixation in mouse tumors induced by 3-methylcholanthrene. *J. Natl. Cancer Inst.* 47, 1255-1265.
216. Willerson JT, Asofsky R, Barth WF (1969) Experimental murine amyloid. IV amyloidosis and immunoglobulins. *J. Immunol.* 103, 741-749.
217. Williams NM, Timoney PJ (1994) Variation in susceptibility of ten mouse strains to infection with a strain of *Ehrlichia risticii*. *J. Comp. Pathol.* 110, 137-143.
218. Willott JF, Aitkin LM, McFadden SL (1993) Plasticity of auditory cortex associated with sensorineural hearing loss in adult C57BL/6J mice. *J. Comp. Neurol.* 329, 402-411.
219. Willott JF, Erway RC, Archer JR, Harrison DE (1995) Genetics of age-related hearing loss in mice. II. Strain differences and effects of caloric restriction on cochlear pathology and evoked response thresholds. *Hearing Research* 88, 143-155.
220. Wimer RE, Wimer CC, Roderick TH (1969) Genetic variability in forebrain structures between inbred strains of mice. *Brain Res.* 16, 257-264.
221. Wragg LE, Speirs RS (1952) Strain and sex differences in response of inbred mice to adrenal cortical hormones. *Proc. Soc. Exp. Biol. Med.* 80, 680-684.
222. Wysocki CJ, Whitney G, Tucker D (1977) Specific anosmia in the laboratory mouse. *Behav. Genet.* 7, 171-188.
223. Xidieh CF, Singer-Vermes LM, Calich VLG, Burger E (1994) Plasma amylase levels as a marker of disease severity in an isogenic murine model of paracoccidioidomycosis. *Journal of Medical and Veterinary Mycology* 32, 37-45.
224. Yamamoto Y, Saito H, Setogawa T, Tomioka H (1991) Sex differences in host resistance to *Mycobacterium marinum* infection in mice. *Infect. Immun.* 59, 4089-4096.
225. Yang GM, Kitagawa K, Matsushita K, Mabuchi T, Yagita Y, Yanagihara T, Matsumoto M (1997) C57BL/6 strain is most susceptible to cerebral ischemia following bilateral common carotid occlusion among seven mouse strains: Selective neuronal death in the murine transient forebrain ischemia. *Brain Res.* 752, 209-218.
226. Young CR, Deacon NJ, Ebringer A, Davis DAL (1976) Genetic control of the immune response to ferritin in mice. *J. Immunogenet.* 3, 199-205.
227. Zarrow MX, Christensen CM, Eleftheriou BC (1971) Strain differences in the ovulatory response of immature mice to PMS and to the pheromonal facilitation of PMS-induced ovulation. *Biol. Reprod.* 4, 52-56.
228. Zelentsov AG (1974) Susceptibility of inbred mice to helminths. II. Development of *Opisthorchis felinus* in A/He, CBA/Lac, CC57W/Mv, C57BL/6J, DBA/2J and SWR/J mice [in Russian]. *Med. Parazit (Mosk.)*, 43, 95-98.
229. Zhang LY, Levitt RC, Kleeberger SR (1995) Differential susceptibility to ozone-induced airways hyperreactivity in inbred strains of mice. *Experimental Lung Research* 21, 503-518.

Contact us

North America 800.793.7287 EU and Asia envigo.com/contactus info@envigo.com



Envigo RMS Division, 8520 Allison Pointe Blvd., Suite 400, Indianapolis, IN 46250, United States