



Pink1 knockout rat - Park6

Model	Pink1 knockout rat - Park6
Strain	HsdSage: LE- <i>Pink1^{tm1Sage}</i>
Location	U.S.
Availability	Live colony

Characteristics/husbandry

- + Homozygous knockout rats exhibit complete loss of target protein
- + Approximately 30% of Pink1 knockout rats display a hindlimb-dragging phenotype at 5 months of age
- + Pink1 knockout rats show increased hindlimb fatigue at 7 weeks of age
- + Pink1 knockout rats show increased number of hindlimb foot slips at 5 and 9 weeks of age as assessed by tapered balance beam
- + Preliminary reports have suggested Pink1 knockout rats show a ~50% reduction in dopaminergic neurons in the substantia nigra at 8 months of age
- + Background strain: Long Evans Hooded

Zygoty genotype

- + Homozygous

Research use

- + Parkinson's disease
- + Stress-induced neurological dysfunction

Origin

The Pink1 (Park6) KO rat model was originally created at SAGE Labs, Inc. in St. Louis, MO and distributed out of the Boyertown, PA facility. The line continues to be maintained through the original SAGE Labs animal inventory acquired by Envigo.

Description

Developed in collaboration with The Michael J. Fox Foundation, this model contains a deletion of the Pink1 (PTEN-induced putative kinase 1) gene, encoding for a serine/threonine protein kinase. Mutations in Pink1 are implicated in early-onset Parkinson's disease. Pink1 knockout rats show both motor impairments and dopaminergic cell loss, making this a useful model of Parkinson's disease.

Pink1 protein kinase localizes to the mitochondria and is thought to protect cells from stress-induced mitochondrial dysfunction. Mutations within this gene result in one form of autosomal recessive early-onset Parkinson's.

In humans, loss of function of Park7 leads to a form of early-onset Parkinson's disease. This occurs due to the role Park7 plays in protecting neurons from oxidative stress and cell death, making this an ideal model for the study of Parkinson's disease.

Figure 1: Increased fatigue in Pink1 knockout rats. Pink1 homozygous and heterozygous knockout rats display decreased movement in an open field after hindlimb fatigue challenge, indicative of increased hindlimb fatigue.






Figure 2: Decreased performance on tapered balance beam in Pink1 knockout rats. Pink1 homozygous and heterozygous knockout rats display increased hindlimb foot slips on a tapered balance beam. No differences were seen in forelimb foot slips.





Figure 3: A graph showing the correlation between the age and weight of Pink1 knockout rats.



Citations

29462410 Garner, Jennifer M; Chambers, Jonathan; Barnes, Abigail K; Datta, Subimal; Changes in Brain-Derived Neurotrophic Factor Expression Influence Sleep-Wake Activity and Homeostatic Regulation of Rapid Eye Movement Sleep. *Sleep* Vol.41, 2018

28576309 Geist, Phillip A; Dulka, Brooke N; Barnes, Abigail; Totty, Michael; Datta, Subimal; BDNF heterozygosity is associated with memory deficits and alterations in cortical and hippocampal EEG power. *Behavioural brain research* Vol.332, 2017

25445195 Gururajan, A; Hill, R A; van den Buuse, M; Brain-derived neurotrophic factor heterozygous mutant rats show selective cognitive changes and vulnerability to chronic corticosterone treatment. *Neuroscience* Vol.284, 2015

26586578 Harris, A P; Lennen, R J; Brydges, N M; Jansen, M A; Pernet, C R; Whalley, H C; Marshall, I; Baker, S; Basso, A M; Day, M; Holmes, M C; Hall, J; The role of brain-derived neurotrophic factor in learned fear processing: an awake rat fMRI study. *Genes, brain, and behavior* Vol.15, 2016

26289587 Liu, Miao; Kay, Jarren C; Shen, Shanwei; Qiao, Li-Ya; Endogenous BDNF augments NMDA receptor phosphorylation in the spinal cord via PLC γ , PKC, and PI3K/Akt pathways during colitis. *Journal of neuroinflammation* Vol.12, 2015

30548775 Martis, Lena-Sophie; Wiborg, Ove; Holmes, Megan C; Harris, Anjanette P; BDNF+/- rats exhibit depressive phenotype and altered expression of genes relevant in mood disorders. *Genes, brain, and behavior* Vol.18, 2019

26506052 Ren, Q; Ma, M; Yang, C; Zhang, J-C; Yao, W; Hashimoto, K; BDNF-TrkB signaling in the nucleus accumbens shell of mice has key role in methamphetamine withdrawal symptoms. *Translational psychiatry* Vol.5, 2015

30855519 Sapio, Matthew R; Iadarola, Michael J; LaPaglia, Danielle M; Lehky, Tanya; Thurm, Audrey E; Danley, Kristen M; Fuhr, Shannon R; Lee, Mark D; Huey, Amanda E; Sharp, Stephen J; Tsao, Jack W; Yanovski, Jack A; Mannes, Andrew J; Han, Joan C; Haploinsufficiency of the brain-derived neurotrophic factor gene is associated with reduced pain sensitivity. *Pain* Vol.160, 2019

26419294 Shirayama, Yukihiko; Yang, Chun; Zhang, Ji-chun; Ren, Qian; Yao, Wei; Hashimoto, Kenji; Alterations in brain-derived neurotrophic factor (BDNF) and its precursor proBDNF in the brain regions of a learned helplessness rat model and the antidepressant effects of a TrkB agonist and antagonist. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* Vol.25, 2015

23583595 St Laurent, Robyn; Helm, Samuel R; Glenn, Melissa J; Reduced cocaine-seeking behavior in heterozygous BDNF knockout rats. *Neuroscience letters* Vol.544, 2013

26687970 Xia, Chunmei; Shen, Shanwei; Hashmi, Fiza; Qiao, Li-Ya; Colitis-induced bladder afferent neuronal activation is regulated by BDNF through PLC β pathway. *Experimental neurology* Vol.285, 2016

27094192 Yang, Bangkun; Yang, Chun; Ren, Qian; Zhang, Ji-Chun; Chen, Qian-Xue; Shirayama, Yukihiko; Hashimoto, Kenji; Regional differences in the expression of brain-derived neurotrophic factor (BDNF) pro-peptide, proBDNF and preproBDNF in the brain confer stress resilience. *European archives of psychiatry and clinical neuroscience* Vol.266, 2016

Contact us

North America 800.793.7287 gemsorders@envigo.com
 Envigo, 8520 Allison Pointe Blvd., Suite 400, Indianapolis, IN 46250, United States

