



Epidermoid Carcinoma Xenograft Tumor Model

MODEL	NOMENCLATURE	HAIR	T CELLS	B CELLS	NK CELLS
Athymic Nude Mouse	Hsd:Athymic Nude-Foxn1 ^{nu}	No	Nonfunctional	Functional	Functional

Model

The athymic nude mouse has an autosomal recessive mutation on *nu* locus on chromosome 11. The hairless model is T-cell deficient and accepts xenograft transplantation.


Cell Line

Human A253 cells sourced from ATCC® (Number: HTB-41™) were implanted into cohorts of athymic nude mice. Female mice at approximately 8 weeks of age were implanted with 1.0e7, 5.0e6, or 1.0e6 cells with GFR Matrigel (1:1 dilution) into the subcutaneous space of the right flank.

Tumor Growth *in vivo*

The mice were maintained in a barrier under controlled environmental conditions. The mice consumed Teklad Global Rodent Diet 2914 (14% protein). Body weights were taken and tumor measurements were assessed with a caliper twice per week.

Tumor Growth Rate for A253 Cells Inoculated into Female Athymic Nude Mice



Data shown as mean values; N=10 per cohort
Tumor growth study services conducted by Covance, Inc.

Contact us

North America 800.793.7287 EU and Asia envigo.com/contactus info@envigo.com